

Supplemental Material

Diagnosis of photochemical ozone production rates and limiting factors in continental outflow air masses reaching Fukue Island, Japan: ozone-control implications

**Yugo Kanaya^{1*}, Hiroshi Tanimoto², Yoko Yokouchi², Fumikazu Taketani¹,
Yuichi Komazaki¹, Hitoshi Irie^{1,3}, Hisahiro Takashima^{1,4}, Xiaole Pan^{1,5},
Susumu Nozoe^{2,6}, Satoshi Inomata²**

¹*Department of Environmental Geochemical Cycle Research, Japan Agency for Marine-Earth Science and Technology, 3173-25 Showa-machi, Kanazawa-ku, Yokohama, Japan*

²*National Institute for Environmental Studies, Tsukuba, Japan*

³*Now at Chiba University, Chiba, Japan*

⁴*Now at Fukuoka University, Fukuoka, Japan*

⁵*Now at Kyushu University, Fukuoka, Japan*

⁶*Now at National Museum of Emerging Science and Innovation, Japan*

The monthly average ozone concentrations at Fukue Island are tabulated in Table S1.

Table S1. Monthly average ozone concentrations (ppb) at Fukue Island

	2009	2010	2011	2012	2013	2014
Jan		41.9	41.0	40.1	47.6	43.9
Feb		44.4	51.7	41.7	47.9	47.5
Mar	51.5	49.0	56.2	54.4	57.5	53.0
Apr	68.9	55.2	61.8	57.5	61.4	61.6
May	60.7	56.9	48.6	60.7	61.9	61.6
Jun	43.0	39.5	35.5	42.8	39.7	45.7
Jul	23.4	30.3	22.1	24.7	22.7	
Aug	29.1		26.5	21.5	29.9	
Sep	43.7		39.5	40.4	46.2	
Oct	51.7	49.7	48.3	49.6	45.0	
Nov	40.3	47.5	39.3	42.9	45.7	
Dec	37.9	40.5	37.9	38.5	42.1	