

Mass concentrations and carbonaceous compositions of PM_{0.1}, PM_{2.5}, and PM₁₀ at urban locations in Hanoi, Vietnam

Nguyen Thi Thu Thuy^{1,2}, Nghiem Trung Dung^{1,*}, Kazuhiko Sekiguchi^{3,*}, Ly Bich Thuy¹, Nguyen Thi Thu Hien¹, Ryosuke Yamaguchi³

¹*School of Environmental Science and Technology, Hanoi University of Science and Technology, 1 Dai Co Viet, Hanoi, Vietnam.*

²*Faculty of International Training, Thai Nguyen University of Technology, 3/2 Road, Thai Nguyen, Vietnam*

³*Graduate School of Science and Engineering, Saitama University, 225 Shimo-Okubo, Sakura, Saitama, Japan*

Supplementary Material

Table S1. Summary of sampling campaign and meteorological conditions

Set	Sampling site	Season	Period (2015)	Number of samples			T (°C)	RH (%)	Ws (m/s)	Wd (°)	P (hPa)	R (w/m ²)	Prd
				PM _{0.1}	PM _{2.5}	PM ₁₀							
1	HUST	Wet	Aug 4 th – Aug 20 th	17 (3)	16 (3)	16 (3)	30.99	74.88	1.59	171.36	1004	195.12	8
2	Vinacomin	Wet	Aug 22 th – Aug 28 th	7 (3)	7 (3)	7 (3)	31.85	70.00	1.42	186.04	1003	204.24	2
3	Vinacomin	Dry	Oct 13 th – Oct 19 th	7 (3)	7(3)	7 (3)	26.72	67.00	0.95	197.07	1014	179.90	0
4	HUST	Dry	Nov 4 th – Dec 30 th	56 (7)	38 (7)	39 (7)	21.70	77.95	1.13	146.26	1017	81.23	25

T, RH, Ws, Wd, P, R, Prd denote the temperature, relative humidity, wind speed, wind direction, surface pressure, the radiation and the number of rainy days in the sampling period, respectively; the value in the brackets denotes the blank samples. The concentrations in this study were calculated at 25° C and 1 atm.