



Near-Surface Vertical Profiles of Urban Roadside NO_x and Fine Particles

Yoshihiro Nakashima¹, Charlotte E. Jones², Wakana Yamanobe³, Yoshizumi Kajii²

¹ Faculty of Agriculture, Tokyo University of Agriculture and Technology, 3-8-1 Harumi-cho, Fuchu, Tokyo, Japan

² Graduate School of Global Environmental Studies, Kyoto University, Yoshida-Honmachi, Sakyo-ku, Kyoto, Japan

³ Aprica Children's Products Japan Co. Ltd, 2-6-12 Ginza, Chuo-ku, Tokyo, Japan

ABSTRACT

This paper presents daytime vertical profiles of the NO_x concentration, and the number concentration and size distribution of fine particles near a major road in urban Tokyo during spring 2011. No significant height dependence was observed in the NO_x concentration, presumably due to rapid diffusion. In contrast, the number concentration of particles under 0.5 μm diameter demonstrated an exponential decrease with increasing height above ground level. Vertical profiles derived from this study differ from those presented in previous studies, however, these differences may potentially be explained by the different tailpipe positions of Japanese and US heavy vehicles, as well as the meteorological conditions. This study demonstrates that in Tokyo, the fine particle concentration at 0.5 m above ground level was about 2.9 times higher than that at 2.0 m. The higher fine particle concentration immediately above ground level implies that children may be at greater risk of experiencing pollutant-related respiratory symptoms than adults.

Keywords: Vertical concentration profile; High density traffic road in urban area; Fine particle.

INTRODUCTION

Exhaust from heavy vehicles contributes one of the major anthropogenic source of the gaseous pollutants CO, NO_x and NMHCs, as well as fine particles. Fine particles in particular are known to cause adverse health effects on humans and a relationship between exposure to high concentrations of fine particles and respiratory morbidity has been suggested (Peden, 2005; Li *et al.*, 2008; Riedl 2008). Recently, an immune response has been observed in humans following exposure to ultrafine particles with a diameter of less than 0.1 μm, (Li *et al.*, 2009, 2010). Several epidemiological studies into respiratory symptoms caused by inhaling anthropogenic fine particles have demonstrated that the prevalence of asthma may be related to regional motor vehicle traffic density and residential proximity to freeways (Holguin 2008; Salam *et al.*, 2008; Patel and Miller, 2009), while other studies reported that there was no evidence to suggest such a relationship (Waldron *et al.*, 1995; Livingstone *et al.*, 1996). To evaluate the correlation between vehicular emissions of fine particles and any potential health impacts, detailed measurements of the concentration and spatial distribution of fine and ultrafine

particles are essential.

The spatial distribution of fine particles has been reported for highway and freeway locations (Zhu *et al.*, 2002; Zhu and Hinds 2005; Zhu *et al.*, 2006; Pohjola *et al.*, 2007; Hagler *et al.*, 2009; Hu *et al.*, 2009; He and Dhaniyala 2012). It has been reported that the horizontal distribution of particle number concentration shows an exponential decay with increasing distance from a highway, in the absence of any obstructions (Zhu *et al.*, 2002). In contrast, the vertical distribution of particles does not indicate exponential decay with increasing distance from the emission source, although the maximum concentration typically occurs close to ground level (< 4 m). These studies were performed in areas surrounding a highway or freeway. However, it is arguably more important to carry out these measurements in the vicinity of high traffic density roads within urban areas.

This study focuses on determining the vertical profiles of the NO_x concentration and number concentration of fine particles within a busy urban street in central Tokyo. The concentrations of NO_x and fine particles were measured at heights from 0.5 to 2.0 m above ground level, since this corresponds to the region where humans will be most directly affected by exposure to vehicular emissions. In theory, young children and babies transported via a stroller may be the most affected by air pollutants near the surface, due to both their relatively low height above ground level and undeveloped respiratory system. The relationship between road traffic and adverse respiratory effects in children has been studied previously (Weiland *et al.*, 1994; Oosterlee *et al.*,

* Corresponding author.

Tel.: +81-42-367-5258; Fax: +81-42-367-5258

E-mail address: nakasima@cc.tuat.ac.jp

1996; Ciccone *et al.*, 1998), and differences in the adverse respiratory effects of fine particles on children and adults have been reported (Oosterlee *et al.*, 1996)

METHODS

Field measurements were conducted in urban Tokyo on April 4th, 2011. Fig. 1 shows the location of the measurement site, which was near to the intersection of two main roads, oriented in the direction of Northeast to Southwest (Route-15) and Northwest to Southeast (Route-304). At a distance of about 300 m from the site, two local roads run parallel to Route-15 in the direction of Northwest (Route-405) and Southeast (Route-316), respectively. In addition, an elevated highway surrounds the measurement site, at a distance of 300–600 m. The area is relatively flat, with tall buildings (6–12 stories) located alongside all roads near to the site. Due to the built up nature of the area, particle emissions from the road adjacent to the site were expected to dominate the measurements, while contributions from other roads in the vicinity are thought to be small. The measurements were carried out between 15:00–16:00 Japanese local time - corresponding to a period of high road usage, with respect to both vehicles and pedestrians. Traffic data for the local roads around the site was obtained from the Ministry of Land, Infrastructure, Transport and Tourism. According to the data, the traffic volume for vehicles in the daytime (7:00–19:00) is approximately 31,000 vehicles per day, with heavy duty vehicles accounting for 16% of the total. The average vehicle speed on the road in typical, congested conditions was 12 km/h.

The size distribution and number concentration of fine particles were measured using an optical particle counter (OPC, model 1.109, Grimm). A laser diode was employed

to detect and count particulates. The size distribution was set from 0.25 to 3.0 μm . Particles of less than 0.25 μm in diameter cannot be detected by the instrument, since these particles are outside of the range for detection. The OPC instrument continuously sampled air at 1.2 L/min and the concentration of each particle size range was measured every 6 seconds. The air was supplied to the instrument via PFA tube (6.35 mm outer diameter, 3 m length) and the inlet of the tube was fixed to a moveable rod, such that the vertical height of the inlet from the ground could be changed periodically. The inlet height was alternated between 0.25, 0.5, 1.0, 1.5 and 2.0 m above ground level. The inlet and OPC were deployed about 1 m away from the intersection. The accumulation time was set to 2 minutes for each measurement height, with an interval of 1 minute following each inlet height adjustment. The full vertical profile measurements were carried out 3 times and the averaged particle number concentration was derived for each height. In addition to the particle measurements, the vertical profile of nitrogen oxides (NO and NO₂) was measured by O₃ chemiluminescence (Model 42i-TL, Thermo Electron) simultaneously via the same inlet system. Unfortunately, meteorological data including ambient temperature, wind speed and direction were not measured due to instrument difficulties. However, it is assumed that the difference of the concentration of the trace species coming from upwind and downwind is small since the roadside are surrounded by the tall buildings.

RESULTS AND DISCUSSIONs

Vertical profiles of the concentrations of NO and NO₂ are shown in Fig. 2. No height dependence was observed in the NO concentration, while a slight height dependence was observed for NO₂, within the standard deviation of the




Fig. 1. An aerial photograph of the measurement site and the surrounding terrain.




Fig. 2. Vertical profiles of the concentration of NO (red) and NO₂ (blue). Plots of NO₂ are shifted to 0.1 m higher in this figure. Error bars indicate one standard deviation.

measurement. Since gaseous species such as NO and NO₂ can readily diffuse within the atmosphere, no strong vertical variation was observed for NO and NO₂ within the range of sampling heights tested. The vertical profiles of pollutant gases have been simulated for CO (Johnson *et al.*, 1973). In scenarios where roadside emission sources are surrounded by tall buildings, the so called “street canyon”, emitted gases are thought to be mixed by helical air circulation. The vertical profiles of pollutant gases for a height z may be derived as follows;

$$C(z) = C(0) \times \frac{x+2}{[(x^2+z^2)^{1/2}+2]} \quad (1)$$

where x stands for the horizontal distance between the emission source and measurement point (2.0 m in the present study), and $C(0)$ is the surface concentration of the pollutant. Applying the Eq. (1) to the present measurement, the ratio of the concentration between the heights of 0.25 and 2.0 m were 0.83. Since the standard deviation of the concentration for each sampling height is large, no height dependence of the concentration simulated by Eq. (1) was observed clearly.

Fig. 3 shows the size distribution of fine particles for each sampling height. Whilst there was virtually no significant height dependence observed in the NO and NO₂ mixing ratios, in contrast, the number concentration of fine particles did demonstrate a degree of height dependence. For all sampling heights, the maximum number concentration was observed for particles of size 0.25–0.30 μm , and the number concentration exponentially decreased with increasing particle size. Fig. 3 indicates that the particle size distribution is broadly uniform throughout all sampling heights, suggesting that there is no unique formation or loss process of the particles occurring at any individual sampling heights. However, the overall number concentration was seen to

decrease with increasing sampling height.

Fig. 4 shows the vertical profiles of the number concentration of particles for each size range. Particles of diameter larger than 0.5 μm showed vertical concentration profiles which were almost uniform (within one standard deviation), while a clear height dependence was observed in the concentration of the smaller particles. The concentration of particles within each size range decreased with increasing height, resulting in apparently exponential vertical profiles of the number concentration. Vertical profiles of the number concentration of fine particles have previously been measured near a highway in September 2009 in Liverpool, New York (He and Dhaniyala, 2012) and close to a freeway in Los Angeles in July 2001 (Zhu and Hinds, 2005). The results from these studies indicate that the maximum particle concentration occurred at a height of 3.4 m (He and Dhaniyala, 2012) and 3.0 m (Zhu and Hinds, 2005). He and Dhaniyala (2012) have explained their results by the position of the tailpipe in heavy duty vehicles and the generation of a thermal plume. Zhu and Hinds, on the other hands, have not explained for the measured vertical profiles but there is the evidence that the ground level of measurement site was ~ 4.5 m lower than the Freeway. In the present study, the tailpipe of heavy vehicles in Japan is located in almost exactly the same position as that of normal commuter gasoline vehicles, ~ 0.5 m above ground level. In addition, the measurements for this study were carried out in late afternoon in April, and as such the ground was not significantly heated since the sunlight was blocked by the tall buildings, which would reduce any thermal plume effect. Ground level of the sampling site is almost same as that of the roadside. These differing conditions are consistent with the differences in the vertical profiles of fine particles measured in this work and the previous studies. Since different vertical profiles of roadside particulates




Fig. 3. Size distribution of the fine particle for each sampling height.




Fig. 4. Vertical profiles of number concentration of fine particle for each particle diameter. Errors bar indicate one standard deviation.

have been observed in different locations, further studies to determine the dependence of the particle number concentration and size distribution on seasonal changes, as well as on the position of the tailpipe are necessary.

For each particle size, the ratio between the standard number concentration at each height and the concentration at 2.0 m was calculated. For particles smaller than 0.5 μm diameter, the ratio of the number concentration increased exponentially as the sampling height decreased. The maximum ratio was 2.9 for the 0.25–0.30 μm size range. This indicates that the level of exposure to fine particles, and therefore the potential for related adverse health effects, may be reduced with increasing height above ground level. In light of this, it might be anticipated that children would be more susceptible to health issues related to roadside fine particle levels compared to adults. Health implications for children living near to areas with high traffic density have been researched previously (Weiland *et al.*, 1994; Oosterlee *et al.*, 1996; Ciccone *et al.*, 1998). Ciccone *et al.* (1998),

investigated chronic respiratory symptoms in children under fifteen years of age and compared them with those of adults living in similar conditions. They concluded that living in the vicinity of busy streets with high traffic flow increased the risk of developing chronic respiratory symptoms in children, while there was no significant correlation in adults. Although they did not offer an explanation for the increased risk in children, fine particles are thought to be a contributing factor.

Based on the results of this study, it is implied the importance of the reduction of the number concentration of fine particle at the lower height. It should be noted that ultrafine particles were not observed in this study, as ambient concentrations were below the instrument detection limit. Based upon the observed vertical profile of fine particles, it is probable that the same height dependence of the number concentration of ultrafine particles will be observed. Recently, a relationship between exposure to ultrafine particles and adverse health effects has been

confirmed (Oberdörster et al., 1994; Penttinen et al., 2001; Li et al., 2010). As such, in future studies to assess the impact of particulate emissions on human health, vertical profile measurements for both fine and ultrafine particles would be desirable.

CONCLUSIONS

Vertical profile measurements of the number concentration of fine particles in an urban street were carried out during the afternoon in spring 2011. The results obtained imply that the risk of developing chronic respiratory symptoms may be greater for children compared to adults. The exponential decay of the particle number concentration with increasing height above ground level is thought to be the result of the low position of tailpipes on Japanese heavy vehicles and the relatively cold road surface. To assess the risk of developing particulate-related respiratory diseases as a result of exposure to vehicular emissions, both the horizontal and vertical distribution of the fine particle concentration should be monitored. In summertime, strong thermal plumes are created in urban areas due to the heat-island phenomenon. As such, it is anticipated that the vertical profile will show seasonal variations. Thus the vertical profile measurements should be repeated in different seasons, in order to achieve a more detailed assessment of typical yearly exposure levels. In addition, in future studies, detailed measurements including ultrafine particles, as well as fine particles should be carried out.

ACKNOWLEDGMENTS

This work was carried out in collaboration with Aprica Children's Products Japan. We would like to thank Chiyo Ohno, Akira Ida and Shungo Kato for their assistance with these measurements.

REFERENCES

- Ciccone, G., Forastiere, F., Agabiti, N., Biggeri, A., Bisanti, L., Chellini, E., Corbo, G., Dell'Orco, V., Dalmaso, P., Volante, T., F., Galassi, C., Piffer, S., Renzoni, E., Rusconi, F., Sestini, P., Viegi, G. and SIDRIA collaborative group (1998). Road Traffic and Adverse Respiratory Effects in Children. *Occup. Environ. Med.* 55(11): 771–778.
- Hagler, G.S.W., Baldauf, R.W., Thoma, E.D., Long, T.R., Snow, R.F., Kinsey, J.S., Oudejans, L. and Gullett, B.K. (2009). Ultrafine Particles near a Major Roadway in Raleigh, North Carolina: Downwind Attenuation and Correlation with Traffic Related Pollutants. *Atmos. Environ.* 43: 1229–1234.
- He, M. and Dhaniyala, S. (2012). Vertical and Horizontal Concentration Distributions of Ultrafine Particles near a Highway. *Atmos. Environ.* 46: 225–236.
- Holguin, F. (2008). Traffic, Outdoor Air Pollution, and Asthma. *Immunol. Allergy Clin. North Am.* 208: 577–588.
- Hu, S., Fruin, S., Kozawa, K., Mara, S., Paulson, S.E. and Winer, A.M. (2009). A Wide Area of Air Pollutant Impact Downwind of a Freeway during Pre-Sunrise Hours. *Atmos. Environ.* 43: 2541–2549.
- Johnson, W.B., Ludwig, F.L., Dabberdt, W.F. and Allen, R.J. (1973). An Urban Diffusion Simulation Model for Carbon Monoxide. *J. Air Pollut. Contr. Assoc.* 23: 490–498.
- Li, N., Xia, T. and Nel, A.E. (2008). The Role of Oxidative Stress in Ambient Particulate Matter-Induced Lung Diseases and its Implications in the Toxicity of Engineered Nanoparticles. *Free Radical Biol. Med.* 44: 1689–1699.
- Li, N., Wang, M., Bramble, L.A., Schmits, D.A., Schauer, J.J., Sioutas, C., Harkema, J.R. and Nel A.E. (2009). The Adjuvant Effect of Ambient Particulate Matter is Closely Reflected by the particulate oxidation potential. *Environ. Health Perspect.* 117: 1116–1123.
- Li, N., Harkema, J.R., Lewandowski, R.P., Wang, M., Bramble, L.A., Gookin, G.R., Ning, Z., Kleinman, M.T., Sioutas, C. and Nel, A.E. (2010). Ambient Ultrafine Particles Provide a Strong Adjuvant Effect in the Secondary Immune Response: Implication for Traffic-Related Asthma Flares. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 299: L374–L383.
- Livingstone A.E., Shaddick, G., Grundy, C. and Elliot, P. (1996). Do People Living near Inner City Main Roads Have More Asthma Needing Treatment? Case Control Study. *BMJ* 312: 676–677.
- Oberdörster, G., Ferin, J. and Lehnert, B.E. (1994). Correlation between Particle Size, in Vivo Particle Persistence, and Lung Injury. *Environ. Health Perspect.* 102: 173–179.
- Oosterlee, A., Drijver, M., Leuret, E. and Brunekreef, B. (1996). Chronic Respiratory Symptoms in Children and Adults Living Along Streets with High Traffic Density. *Occup. Environ. Med.* 53: 241–247.
- Patel, M.M. and Miller R.L. (2009). Air Pollution and Childhood Asthma: Recent Advanced and Future Directions. *Curr. Opin. Pediatr.* 21: 235–242.
- Peden, D.B. (2005). The Epidemiology and Genetics of Asthma Risk Associated with Air Pollution. *J. Allergy Clin. Immunol.* 115: 213–219.
- Penttinen, P., Timonen, K.L., Tittanen, P., Mirme, A., Ruuskanen, J. and Pekkanen, J. (2001). Ultrafine Particles in Urban Air and Respiratory Health among Adult Asthmatics. *Eur. Respir. J.* 17: 428–435.
- Pohjola, M.A., Pirjola, L., Karppinen, A., Harkonen, J., Korhonen, H., Hussein, T., Ketzel, M. and Kukkonen, J. (2007). Evaluation and Modeling of the Size Fractionated Aerosol Particle Number Concentration Measurements nearby a Major Road in Helsinki - part I: Modeling Results within the LIPIKA Project. *Atmos. Chem. Phys.* 7: 4065–4080.
- Riedl, M.A. (2008). The Effect of Air Pollution on Asthma and Allergy. *Curr. Allergy Asthma Rep.* 8: 139–146.
- Salam, M.T., Islam, T. and Gilliland, F.D. (2008). Recent Evidence for Adverse Effects of Residential Proximity to Traffic Sources on Asthma. *Curr. Opin. Pulm. Med.* 14: 3–8.
- Waldron, G., Pottle, B. and Dodd, J. (1995). Asthma and the Motorways- one District's Experience. *J. Public*

- Health Med.* 17: 85–89.
- Weiland, S.K., Mundt, K.A., Rückmann, A. and Keil, U. (1994). Self Reported Wheezing and Allergic Rhinitis in Children and Traffic Density on Street of Residence. *Ann. Epidemiol.* 4: 243–247.
- Zhu, Y., Hinds, W.C., Kim, S. and Sioutas, C. (2002). Concentration and Size Distribution of Ultrafine Particles near a Major Highway. *J. Air Waste Manage. Assoc.* 52: 1032–1042.
- Zhu, Y. and Hinds, W.C. (2005). Predicting Particle Number Concentrations near a Highway Based on Vertical Concentration Profile. *Atmos. Environ.* 39: 1557–1566.
- Zhu, Y., Kuhn, T., Mayo, P. and Hinds, W.C. (2006). Comparison of Daytime and Nighttime Concentration Profiles and Size Distributions of Ultrafine Particles near a Major Highway. *Environ. Sci. Technol.* 40: 2531–2536.

Received for review, August 24, 2013

Accepted, January 1, 2014